

MIGRAINE AND HEADACHE TREATMENTS (Part 1 of 3)

Generic	Strength	Brand	Form	Dose
ANALGESIC + BARBITURATE				
acetaminophen	325mg	—	℞ caps, tabs	Adults: 1–2 tabs or caps every 4hrs as needed; max 6/day Children: Not established.
butalbital	50mg			
caffeine	40mg			
acetaminophen	300mg	Fioricet	℞ caps	Adults: 1–2 caps every 4hrs; max 6/day Children: Not recommended.
butalbital	50mg			
caffeine	40mg			
acetaminophen	325mg	—	℞ tabs	Adults: 1–2 tabs every 4hrs; max 6/day. Children: Not recommended.
butalbital	50mg			
ANALGESIC + BARBITURATE + OPIOID				
acetaminophen	300mg	Fioricet w. Codeine	CIII caps	Adults: 1–2 caps every 4hrs; max 6/day. Children: Not recommended.
butalbital	50mg			
caffeine	40mg			
codeine phosphate	30mg			
ANALGESIC + SALICYLATE				
acetaminophen	250mg	Bayer Migraine Formula	OTC caplets	Adults: 2 caplets with a glass of water; max 2 caplets/24hrs. Children: Not recommended.
aspirin	250mg			
caffeine	65mg			
acetaminophen	250mg	Excedrin Migraine	OTC caplets	Adults: 2 caplets with a glass of water; max 2/day. Children: Consult physician.
aspirin	250mg			
caffeine	65mg			
BARBITURATE + SALICYLATE				
aspirin	325mg	Fiorinal	CIII tabs, caps	Adults: 1–2 tabs or caps every 4hrs as needed; max 6/day. Children: Not recommended.
butalbital	50mg			
caffeine	40mg			
BARBITURATE + SALICYLATE + OPIOID				
aspirin	325mg	Fiorinal w. Codeine	CIII caps	Adults: 1–2 tabs or caps every 4hrs as needed; max 6/day. Children: Not recommended.
butalbital	50mg			
caffeine	40mg			
codeine phosphate	30mg			
ERGOT ALKALOID				
dihydroergotamine mesylate	1mg/mL	D.H.E. 45	℞ IM, IV, or SC inj	Adults: 1mL IV at 1hr intervals; max 2 doses/day. Or, 1mL IM or SC at 1hr intervals; max 3 doses/day. For all: max 6 doses/wk. Not for chronic use. Children: Not recommended.
	4mg/mL	Migranal	℞ nasal spray†	
ergotamine tartrate	1mg	—	℞ tabs	Adults: 2 tabs at onset of attack, then 1 tab every ½hr if needed; max 6 tabs/attack, 10 tabs/wk. Children: Not recommended.
caffeine	100mg			
ergotamine tartrate	2mg	—	℞ rectal supp	Adults: 1 supp rectally at onset of attack, then 1 supp after 1hr if needed; max 2 supps/attack, 5 supps/wk. Children: Not recommended.
caffeine	100mg			
NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)				
diclofenac potassium	50mg	Cambia	℞ buffered pwd for oral soln*	Adults: Mix 1 packet (50mg) with 30–60mL of water only and drink immediately. Not interchangeable with other forms of diclofenac.. Children: Not recommended.

(continued)

MIGRAINE AND HEADACHE TREATMENTS (Part 2 of 3)

Generic	Strength	Brand	Form	Dose
NON-STEROIDAL ANTI-INFLAMMATORY (NSAID) (continued)				
ibuprofen	200mg	Advil Migraine	OTC liquid-filled caps	Adults: Usually 400mg once daily. Children: Not recommended.
	200mg	Motrin Migraine Pain	OTC caplets	Adults: Usually 200–400mg once daily. Children: Not recommended.
SELECTIVE 5-HT_{1B/1D} RECEPTOR AGONIST				
almotriptan (as malate)	6.25mg, 12.5mg	Axert	℞ tabs	Adults: 6.25–12.5mg single dose; may repeat once after 2hrs; max 25mg/24hrs. <i>Hepatic impairment, CrCl 10–30mL/min, or concomitant potent CYP3A4 inhibitors:</i> initially 6.25mg once; max 12.5mg/24hrs. The safety of treating an average of more than 4 migraines over 30 days is not established. Children: Not established.
eletriptan HBr	20mg, 40mg	Relpax	℞ tabs	Adults: 20mg or 40mg once. May repeat once after 2hrs; max 80mg/day. The safety of treating an average of more than 3 headaches in a 30-day period has not been established. Children: Not established.
frovatriptan (as succinate)	2.5mg	Frova	℞ tabs	Adults: 2.5mg with fluids; may repeat once after 2hrs; max 7.5mg/24hrs. The safety of treating an average of more than 4 headaches in a 30-day period has not been established. Children: Not recommended.
naratriptan HCl	1mg, 2.5mg	Amerge	℞ tabs	Adults: 1mg or 2.5mg with fluids; may repeat once after 4hrs; max 5mg/24hrs. The safety of treating, on average, more than 4 headaches in a 30-day period has not been established. <i>Mild-to-moderate renal/hepatic impairment:</i> initially 1mg; max 2.5mg/24hrs. Children: Not recommended.
rizatriptan (as benzoate)	5mg, 10mg	Maxalt	℞ tabs	Adults: Initially 5 or 10mg; may repeat after 2hrs; max 30mg/day. <i>Concomitant propranolol:</i> 5mg; max 3 doses/day. The safety of treating, on average, more than 4 headaches in a 30-day period has not been established. Children: Not recommended.
	5mg, 10mg	Maxalt-MLT	℞ ODT*	
sumatriptan (as succinate)	6mg/0.5mL	Alsuma	℞ soln for SC inj	Adults: One (6mg) dose SC to the lateral thigh or upper arm only. May repeat after 1hr; max two (6mg) doses in 24hrs. Children: Not recommended.
	4mg/0.5mL, 6mg/0.5mL	Imitrex Injection	℞ SC inj	Adults: 6mg SC (may start at lower dose if 6mg not tolerated). May repeat after 1hr; max two 6mg doses in 24hrs. Children: Not recommended.
	5mg/spray, 20mg/spray	Imitrex Nasal Spray	℞ nasal spray	Adults: 5mg, 10mg, or 20mg once. May repeat once after 2hrs; max 40mg/day. The safety of treating an average of more than 4 headaches in a 30-day period has not been established. Children: Not recommended.
	25mg, 50mg, 100mg	Imitrex Tablets	℞ tabs	Adults: 25–100mg once, swallow whole with fluids as soon as possible after migraine onset; may repeat dose at intervals of at least 2hrs, max 200mg/day; or single-dose tablets up to 100mg/day if injection has been used. The safety of treating an average of more than 4 headaches in a 30-day period has not been established. Children: Not recommended.

(continued)

MIGRAINE AND HEADACHE TREATMENTS (Part 3 of 3)

Generic	Strength	Brand	Form	Dose
SELECTIVE 5-HT_{1B/1D} RECEPTOR AGONIST (continued)				
sumatriptan (as succinate) (continued)	11mg	Onzetra Xsail	℞ caps for nasal inh	Adults: 1 nosepiece (11mg) in each nostril (22mg total), using Xsail delivery device. May repeat once after ≥2hrs; max 44mg (4 nosepieces)/day or one dose of Onzetra Xsail and one dose of another sumatriptan product separated by ≥2hrs. The safety of treating an average of more than 4 headaches in a 30-day period has not been established. Children: Not established.
	4mg/0.5mL, 6mg/0.5mL	Sumavel DosePro	℞ soln for SC inj	Adults: <i>Cluster headache:</i> one (6mg) DosePro SC to the abdomen or thigh only. <i>Migraine:</i> may use one (4mg) or (6mg) DosePro SC to the abdomen or thigh only. May repeat after 1hr; max 12mg in 24hrs. Children: Not recommended.
	6.5mg/4hrs	Zecuity	℞ transdermal system	Adults: 1 transdermal system applied to the upper arm or thigh only. Max 2 systems in 24hrs; 2nd system should be applied to a different site no sooner than 2hrs after activation of the 1st system. Children: Not recommended.
	3mg/0.5mL	Zembrace Symtouch	℞ soln for SC inj	Adults: 3mg SC. May repeat after ≥1hr; max 12mg in 24hrs. Children: Not recommended.
zolmitriptan	2.5mg+, 5mg	Zomig	℞ tabs	Adults: Initially 1.25–2.5mg; max recommended single dose: 5mg. If headache returns, may repeat after 2hrs; max 10mg/day. The safety of treating an average of more than 3 headaches in a 30-day period has not been established. <i>Concomitant cimetidine:</i> max single dose: 2.5mg, not to exceed 5mg in 24-hr period. Children: Not recommended.
	5mg/spray	Zomig Nasal Spray	℞ nasal spray	Adults: 5mg once. If headache returns, may repeat once after 2hrs; max 10mg/day. The safety of treating an average of more than 4 headaches in a 30-day period has not been established. Children: Not recommended.
	2.5mg, 5mg	Zomig-ZMT	℞ ODT*	Adults: Initially 1.25–2.5mg; max recommended single dose: 5mg. If headache returns, may repeat after 2hrs; max 10mg/day. The safety of treating an average of more than 3 headaches in a 30-day period has not been established. <i>Concomitant cimetidine:</i> max single dose: 2.5mg, not to exceed 5mg in 24-hr period. Children: Not recommended.
SELECTIVE 5-HT_{1B/1D} RECEPTOR AGONIST + NON-STEROIDAL ANTI-INFLAMMATORY (NSAID)				
sumatriptan succinate/naproxen sodium	10/60mg, 85/500mg	Treximet	℞ tabs	Adults: 1 tab (85/500mg) once; may repeat once after 2hrs; max 2 tabs/day. <i>Mild-to-moderate hepatic impairment:</i> 1 tab (10/60mg)/day. The safety of treating an average of more than 5 migraines in a 30-day period has not been established. Children: <12yrs: Not established. 12–17yrs: 1 tab (10/60mg) once; max 1 tab (85/500mg)/day. The safety of treating an average of more than 2 migraines in a 30-day period has not been established.

NOTES

Key: ODT = orally disintegrating tablets; soln = solution; supp = suppositories; + = scored tablets

*contains phenylalanine; †contains caffeine

Not an inclusive list of medications, official indications and/or dosages. Please see drug monograph at www.eMPR.com, contact company for full drug labeling and/or consult your healthcare provider.

(Rev. 3/2017)