


Classifying Asthma Severity and Initiating Treatment in Children 5 to 11 Years of Age

Assessing severity and initiating therapy in children who are not currently taking long-term control medication

Components of Severity		Classification of Asthma Severity (5 to 11 Years of Age)			
		Intermittent	Persistent		
			Mild	Moderate	Severe
Impairment	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day
	Nighttime awakenings	≤2×/month	3–4×/month	>1×/week but not nightly	Often 7×/week
	Short-acting β ₂ -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not daily	Daily	Several times per day
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited
	Lung function	<ul style="list-style-type: none"> Normal FEV₁ between exacerbations FEV₁ >80% predicted FEV₁/FVC >85% 	<ul style="list-style-type: none"> FEV₁ ≥80% predicted FEV₁/FVC >80% 	<ul style="list-style-type: none"> FEV₁ = 60%–80% predicted FEV₁/FVC = 75%–80% 	<ul style="list-style-type: none"> FEV₁ <60% predicted FEV₁/FVC <75%
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2/year		
		← Consider severity and interval since last exacerbation → Frequency and severity may fluctuate over time for patients in any severity category			
		Relative annual risk of exacerbations may be related to FEV ₁			
Recommended Step for Initiating Therapy		Step 1	Step 2	Step 3, medium-dose ICS option	Step 3, medium-dose ICS option, or Step 4
		and consider short course of oral systemic corticosteroids			
In 2 to 6 weeks, evaluate level of asthma control that is achieved and adjust therapy accordingly.					

EIB = exercise-induced bronchospasm; FEV₁ = forced expiratory volume in 1 second; FVC = forced vital capacity; ICS = inhaled corticosteroid. Adapted from National Asthma Education and Prevention Program. *Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma* 2007. U.S. Department of Health and Human Services. Available at: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf>. Accessed on: November 26, 2012.

Stepwise Approach for Managing Asthma in Children 5 to 11 Years of Age



ICS = inhaled corticosteroid; LABA = inhaled long-acting β₂-agonist; LTRA = leukotriene receptor antagonist; SABA = inhaled short-acting β₂-agonist. Adapted from National Asthma Education and Prevention Program. *Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma* 2007. U.S. Department of Health and Human Services. Available at: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf>. Accessed on: November 26, 2012.

Assessing Asthma Control and Adjusting Therapy in Children 5 to 11 Years of Age

Components of Control		Classification of Asthma Control (5 to 11 Years of Age)		
		Well Controlled	Not Well Controlled	Very Poorly Controlled
Impairment	Symptoms	≤2 days/week but not more than once on each day	>2 days/week or multiple times on ≤2 days/week	Throughout the day
	Nighttime awakenings	≤1×/month	≥2×/month	≥2×/week
	Interference with normal activity	None	Some limitation	Extremely limited
	Short-acting β_2 -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week	Several times per day
	Lung function • FEV ₁ or peak flow • FEV ₁ /FVC	>80% predicted/ personal best >80%	60%–80% predicted/ personal best 75%–80%	<60% predicted/ personal best <75%
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2/year	
		Consider severity and interval since last exacerbation		
	Reduction in lung growth	Evaluation requires long-term follow-up		
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.		
Recommended Action for Treatment		<ul style="list-style-type: none"> • Maintain current step • Regular follow-up every 1 to 6 months • Consider step down if well controlled for at least 3 months 	<ul style="list-style-type: none"> • Step up at least 1 step and • Reevaluate in 2 to 6 weeks • For side effects, consider alternative treatment options 	<ul style="list-style-type: none"> • Consider short course of oral systemic corticosteroids • Step up 1 to 2 steps, and • Reevaluate in 2 weeks • For side effects, consider alternative treatment options

EIB = exercise-induced bronchospasm; FEV₁ = forced expiratory volume in 1 second; FVC = forced vital capacity.

Adapted from National Asthma Education and Prevention Program. *Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma 2007*. U.S. Department of Health and Human Services. Available at: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf>. Accessed on: November 26, 2012.

This Asthma Management Chart is also available at www.eMPR.com.

(Rev. 12/2012)