

ASTHMA MANAGEMENT: ≥12 YEARS OF AGE (Part 1 of 2)

Classifying Asthma Severity and Initiating Treatment

Assessing severity and initiating treatment for patients who are not currently taking long-term control medications

Components of Severity		Classification of Asthma Severity (≥12 Years of Age)				
		Intermittent	Persistent			
			Mild	Moderate	Severe	
Impairment Normal FEV₁/FVC: 8–19yr 85% 20–39yr 80% 40–59yr 75% 60–80yr 70%	Symptoms	≤2 days/week	>2 days/week but not daily	Daily	Throughout the day	
	Nighttime awakenings	≤2×/month	3–4×/month	>1×/week but not nightly	Often 7×/week	
	Short-acting β_2 -agonist use for symptom control (not prevention of EIB)	≤2 days/week	>2 days/week but not daily and not more than 1× on any day	Daily	Several times per day	
	Interference with normal activity	None	Minor limitation	Some limitation	Extremely limited	
	Lung function	<ul style="list-style-type: none"> • Normal FEV₁ between exacerbations • FEV₁ >80% predicted • FEV₁/FVC normal 	<ul style="list-style-type: none"> • FEV₁ >80% predicted • FEV₁/FVC normal 	<ul style="list-style-type: none"> • FEV₁ >60% but <80% predicted • FEV₁/FVC reduced 5% 	<ul style="list-style-type: none"> • FEV₁ <60% predicted • FEV₁/FVC reduced >5% 	
Risk	Exacerbations requiring oral systemic corticosteroids	0–1/year	≥2/year	 Consider severity and interval since last exacerbation		
		 Frequency and severity may fluctuate over time for patients in any severity category				
Recommended Step for Initiating Treatment		Step 1	Step 2	Step 3	Step 4 or 5	
In 2–6 weeks, evaluate level of asthma control that is achieved and adjust therapy accordingly.						

Stepwise Approach for Managing Asthma

Intermittent Asthma	Persistent Asthma: Daily Medication Consult with asthma specialist if Step 4 care or higher is required. Consider consultation at Step 3.			Step 6 Preferred: * High-dose ICS + LABA + oral corticosteroid AND Consider Omalizumab for patients who have allergies	Step up if needed (first, check adherence, environmental control, and comorbid conditions)
Step 1 Preferred: SABA PRN	Step 2 Preferred: Low-dose ICS Alternative: Cromolyn, LTRA, Nedocromil, or Theophylline	Step 3 Preferred: Low-dose ICS + LABA OR Medium-dose ICS Alternative: Low-dose ICS + either LTRA, Theophylline, or Zileuton	Step 4 Preferred: Medium-dose ICS + LABA Alternative: Medium-dose ICS + either LTRA, Theophylline, or Zileuton	Step 5 Preferred: High-dose ICS + LABA AND Consider Omalizumab for patients who have allergies	Assess control
Each Step: Patient education, environmental control, and management of comorbidities Steps 2–4: Consider subcutaneous allergen immunotherapy for patients who have allergic asthma.					
Quick-Relief Medication for All Patients <ul style="list-style-type: none"> • SABA as needed for symptoms. Intensity of treatment depends on severity of symptoms: up to 3 treatments at 20-min intervals as needed. Short course of oral systemic corticosteroids may be needed • Use of SABA >2 days a week for symptom relief (not prevention of EIB) generally indicates inadequate control and the need to step up treatment 					

NOTES

Key: EIB = exercise-induced bronchospasm; FEV₁ = forced expiratory volume in 1 second; FVC = forced vital capacity; ICS = inhaled corticosteroid; LABA = inhaled long-acting β_2 -agonist; LTRA = leukotriene receptor antagonist; SABA = inhaled short-acting β_2 -agonist.

* Preferred therapy is based on Expert Panel Report 2 from 1997.

(continued)

ASTHMA MANAGEMENT: ≥ 12 YEARS OF AGE (Part 2 of 2)

Assessing Asthma Control and Adjusting Therapy

Components of Control		Classification of Asthma Control (≥ 12 Years of Age)				
		Well Controlled	Not Well Controlled	Very Poorly Controlled		
Impairment	Symptoms	≤ 2 days/week	>2 days/week	Throughout the day		
	Nighttime awakenings	$\leq 2 \times$ /month	1–3 \times /week	$\geq 4 \times$ /week		
	Interference with normal activity	None	Some limitation	Extremely limited		
	Short-acting β_2 -agonist use for symptom control (not prevention of EIB)	≤ 2 days/week	>2 days/week	Several times per day		
	FEV ₁ or peak flow	$>80\%$ predicted/ personal best	60%–80% predicted/ personal best	$<60\%$ predicted/ personal best		
	Validated questionnaires*	ATAQ 0 ACQ $\leq 0.75^\dagger$ ACT ≥ 20	1–2 ≥ 1.5 16–19	3–4 N/A ≤ 15		
	Exacerbations requiring oral systemic corticosteroids	0–1/year	$\geq 2/year$			
Risk	Progressive loss of lung function	Evaluation requires long-term follow-up care				
	Treatment-related adverse effects	Medication side effects can vary in intensity from none to very troublesome and worrisome. The level of intensity does not correlate to specific levels of control but should be considered in the overall assessment of risk.				
Recommended Action for Treatment		<ul style="list-style-type: none"> Maintain current step Regular follow-ups every 1–6 months to maintain control Consider step down if well controlled for at least 3 months 	<ul style="list-style-type: none"> Step up 1 step and reevaluate in 2–6 weeks For side effects, consider alternative treatment options 	<ul style="list-style-type: none"> Consider short course of oral systemic corticosteroids Step up 1 to 2 steps and reevaluate in 2 weeks For side effects, consider alternative treatment options 		

NOTES

Key: ACQ = Asthma Control Questionnaire[®]; ACT = Asthma Control Test[™]; ATAQ = Asthma Therapy Assessment Questionnaire[®]; EIB = exercise-induced bronchospasm; FEV₁ = forced expiratory volume in 1 second.

*Questionnaires do not assess lung function or the risk domain.

[†]ACQ values of 0.76–1.4 are indeterminate regarding well-controlled asthma.

REFERENCES

Adapted from National Asthma Education and Prevention Program. *Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma* 2007. U.S. Department of Health and Human Services. Available at: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdn.pdf>. Accessed on: November 26, 2012. (Rev. 7/2013)