

HEPATITIS B DOSING*

Brand	Generic	Usual Dosage
Baraclude	entecavir	Adults: ≥16yrs: Take on empty stomach. Compensated liver disease: nucleoside-naïve: 0.5mg once daily; lamivudine-refractory or known lamivudine or telbivudine resistance mutations: 1mg once daily. Decompensated liver disease: 1mg once daily. Children: <16yrs: Not recommended.
EpiVir-HBV	lamivudine	Adults: 100mg once daily. Children: 2–17yrs: 3mg/kg (max 100mg) once daily; <2yrs: Not recommended.
Hepsera	adefovir	Adults and Children ≥12yrs: 10mg once daily.
Intron A	interferon alfa-2b	Adults: 5million IU daily or 10million IU IM or SC three times weekly for 16 wks. Children: ≥1yr: 3million IU/m ² SC three times weekly for 1 week, then increase to 6million IU/m ² (max 10million IU) 3 times weekly for a total of 16–24 wks; <1yr: Not recommended.
Pegasys	peginterferon alfa-2a	Adults: ≥18yrs: 180mcg SC once weekly for 48 wks. Children: <18yrs: Not recommended.
Tyzeka	telbivudine	Adults: ≥16yrs: 600mg once daily. Children: <16yrs: Not recommended.
Viread	tenofovir	Adults and Children ≥12yrs (≥35kg): 300mg once daily.

NOTES

*For Hepatitis B Adult Renal Dosing Chart or full prescribing information, refer to individual drug monographs found at www.empr.com.

(Rev. 2/2013)