Generic	Brand	Strength	Size	Form	Adult Dose			
ALPHA <sub>2</sub> -AGONISTS								
apraclonidine HCl	lopidine	0.5%	5mL, 10mL	soln <sup>1</sup>	1–2 drops 3 times daily			
brimonidine tartrate	Alphagan P	0.1%, 0.15%	5mL, 10mL, 15mL	soln <sup>2</sup>	1 drop every 8hrs			
ALPHA <sub>2</sub> -AGONIST + BETA-BLOCKER								
brimonidine tartrate/ timolol maleate	Combigan	0.2%/0.5%	5mL, 10mL	soln <sup>1</sup>	1 drop every 12hrs			
BETA-BLOCKERS								
betaxolol HCl	<u> </u>	0.5%	5mL, 10mL, 15mL	soln <sup>1</sup>	1–2 drops twice daily			
	Betoptic S	0.25%	2.5mL, 5mL, 10mL, 15mL	susp <sup>1</sup>	1–2 drops twice daily			
carteolol HCl	_	1%	5mL, 10mL, 15mL	soln <sup>1</sup>	1 drop twice daily			
levobunolol HCl	Betagan	0.5%	5mL, 10mL, 15mL	soln <sup>1</sup>	1–2 drops once or twice daily			
metipranolol HCl	Optipranolol	0.3%	5mL, 10mL	soln <sup>1</sup>	1 drop twice daily			

soln1

soln1

soln1

soln<sup>4</sup>

gelforming soln

sust-rel

caps

soln1

tahs

susp1

soln1

soln4

soln1

soln1

gel1

2.5mL, 5mL, 10mL, susp1

1 drop twice daily

1 drop once daily

500mg twice daily

1 drop 3 times daily

1 drop 3 times daily

1 drop three times daily

2 drops up to 3 times daily

2 drops up to 3–4 times daily or more if needed. *Heavily pigmented irides:* Need higher concentrations.

Apply 1/2 inch ribbon in lower

(continued)

conjunctival sac at bedtime

1 drop twice daily

50-100mg twice to three times daily

1 drop once daily in the morning

Initially 1 drop of 0.25% twice daily; max 1 drop of 0.5% twice daily

5mL, 10mL, 15mL

5ml

60

100

15mL

10ml

90

8ml

10ml

0.2mL

15mL

15mL

4q

15mL, 30mL

5mL, 10mL

2.5mL, 5mL

0.25%, 0.5%

0.25%, 0.5%

0.25%, 0.5%

500ma

1%

2%

25mg, 50mg

1%/0.2%

2%/0.5%

2%/0.5%

1%, 2%, 4%

: 1.5%

3%

4%

0.5%

**Timoptic Ocudose** 0.25%, 0.5%

timolol hemihydrate

timolol maleate

acetazolamide

brinzolamide

dorzolamide HCl

methazolamide

brinzolamide/

brimonidine tartrate

dorzolamide HCI/

timolol maleate

carbachol

pilocarpine HCl

**Betimol** 

Istalol

CARBONIC ANHYDRASE INHIBITORS

Azopt

Trusopt

Neptazane

Simbrinza

Cosopt

CHOLINERGIC (MIOTIC) AGONISTS

Cosopt PF

CARBONIC ANHYDRASE INHIBITOR + ALPHA<sub>2</sub>-AGONIST

CARBONIC ANHYDRASE INHIBITOR + BETA-BLOCKER

Isopto Carbachol

**Isopto Carpine** 

Pilopine HS

Timoptic-XE

Diamox Sequels

**GLAUCOMA TREATMENTS** (Part 1 of 2)

GEAGCOMA INCAMILITIS (I dit 2012)									
Generic	Brand	Strength	Size	Form	Adult Dose				
CHOLINESTERASE INHIBITORS									
ecothiophate iodide	lodide	0.03%, 0.06%, 0.125%, 0.25	5mL		0.03% soln instilled twice daily. If being transferred due to unsatisfactory control with pilocarpine, carbachol, epinephrine, etc, use higher strength soln.				
PROSTAGLANDIN ANALOGS									
bimatoprost	Lumigan	0.01%, 0.03%	2.5mL, 5mL, 7.5mL	soln <sup>1</sup>	1 drop once daily in the evening				
latanoprost	Xalatan	0.005%	2.5mL	soln <sup>1</sup>	1 drop once daily in the evening				
tafluprost	Zioptan	0.0015%	0.3mL	soln <sup>4</sup>	1 drop once daily in the evening				
travoprost	Travatan Z	0.004%	2.5mL, 5mL	soln <sup>3</sup>	1 drop once daily in the evening				
PROSTANOIDS (DO	OCOSANOIDS)								
unoprostone isopropyl	Rescula	0.15%	5mL	soln <sup>1</sup>	1 drop twice daily				
SYMPATHOMIMET	TICS								
dipivefrin HCl	Propine	0.1%	5mL, 10mL, 15mL	soln <sup>1</sup>	1 drop every 12hrs				

**GLAUCOMA TREATMENTS** (Part 2 of 2)

**NOTES** 

<sup>2</sup>Contains purite. <sup>3</sup>Contains *sof* Zia (boric acid, propylene glycol, sorbitol, zinc chloride). <sup>4</sup>Preservative-free.

(Rev. 8/2017)

Not an inclusive list of medications and/or official indications. Please see drug monograph at www.eMPR.com and/or contact company for full drug labeling.

**Key:** soln = solution; sust-rel caps = sustained release capsules; susp = suspension <sup>1</sup>Contains benzalkonium chloride. May be absorbed by soft contact lenses.